Use of fresh stylosanthes (Styosanthes guianensis, CIAT 184) and cassava foliage (Manihot esculenta, Crantz) as a protein source for crossbred pigs

Bounlieng Khoutsavang

Livestock Research Centre, National Agriculture and Forestry Research Institute. Ministry of Agriculture and Forestry, P.O. Box 811, Vientiane, Lao PDR

Acknowledgements

The studies in this thesis were carried out at the Livestock Research Centre, National Agriculture and Forestry Research Institute, Vientiane, Lao PDR. The author gratefully acknowledges the Swedish International Development Co-operation Agency, Department for Research Cooperation (Sida-SAREC) for its financial and material support of this study and I would also like to express my great gratitude to:

Prof. Dr. R. Brian Ogle, Director of the Master of Science course on "Sustainable Livestock Production Systems in the Tropics" my supervisor, for his support, valuable advice and useful guidance to me during my study and I am greatly indebted to him.

Dr. Bounthong Bouahom. Director of the National Agriculture and Forestry Research Institute (NAFRI) for giving me permission to study, facilitation and encouragement

The professors, doctors and lecturers who provided me with useful knowledge during the MSc course

Livestock Research Centre staff for providing facilities and the site for the research work, and also thanks to MSc. Phonepaseuth Phengsavanh for his useful advice and help with the statistical analysis for my research work

My classmates in the MSc course for their contributions and suggestions during my study time.